

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
13 November 2003 (13.11.2003)

PCT

(10) International Publication Number
WO 03/093264 A1

(51) International Patent Classification⁷: **C07D 417/04**,
417/14, A61K 31/433, 31/4439, A61P 25/28

(74) Agent: **BUCHAN, Gavin, MacNicol**; Merck & Co., Inc.,
European Patent Department, Terlings Park, Eastwick
Road, Harlow, Essex CM20 2QR (GB).

(21) International Application Number: **PCT/GB03/01777**

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 24 April 2003 (24.04.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0209995.0 1 May 2002 (01.05.2002) GB

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **MERCK
SHARP & DOHME LIMITED** [GB/GB]; Hertford Road,
Hoddesdon, Hertfordshire EN11 9BU (GB).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): **COLLINS, Ian**,
James [GB/GB]; Terlings Park, Eastwick Road, Harlow,
Essex CM20 2QR (GB). **COOPER, Laura, Catherine**
[GB/GB]; Terlings Park, Eastwick Road, Harlow, Essex
CM20 2QR (GB).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **OXADIAZOLE DERIVATIVES FOR INHIBITION OF GAMMA SECRETASE**



(57) Abstract: Compounds of formula I: wherein X represents a 5-(R-substituted)-1,2,4-oxadiazole-3-yl moiety are inhibitors of gamma-secretase, and hence useful for treating Alzheimer's disease.

WO 03/093264 A1

**OXADIAZOLE DERIVATIVES FOR INHIBITION OF GAMMA
SECRETASE**

The present invention relates to a novel class of compounds, their salts, pharmaceutical compositions comprising them, processes for making them
5 and their use in therapy of the human body. In particular, the invention relates to compounds which modulate the processing of APP by γ -secretase, and hence are useful in the treatment or prevention of Alzheimer's disease.

Alzheimer's disease (AD) is the most prevalent form of dementia.
10 Although primarily a disease of the elderly, affecting up to 10% of the population over the age of 65, AD also affects significant numbers of younger patients with a genetic predisposition. It is a neurodegenerative disorder, clinically characterized by progressive loss of memory and cognitive function, and pathologically characterized by the deposition of
15 extracellular proteinaceous plaques in the cortical and associative brain regions of sufferers. These plaques mainly comprise fibrillar aggregates of β -amyloid peptide ($A\beta$), and although the exact role of the plaques in the onset and progress of AD is not fully understood, it is generally accepted that suppressing or attenuating the secretion of $A\beta$ is a likely means of
20 alleviating or preventing the condition. (See, for example, *ID research alert* 1996 1(2):1-7; *ID research alert* 1997 2(1):1-8; Current Opinion in CPNS Investigational Drugs 1999 1(3):327-332; and Chemistry in Britain, Jan. 2000, 28-31.)

$A\beta$ is a peptide comprising 39-43 amino acid residues, formed by
25 proteolysis of the much larger amyloid precursor protein. The amyloid precursor protein (APP or $A\beta$ PP) has a receptor-like structure with a large ectodomain, a membrane spanning region and a short cytoplasmic tail. Different isoforms of APP result from the alternative splicing of three exons in a single gene and have 695, 751 and 770 amino acids respectively.

30 The $A\beta$ domain encompasses parts of both extra-cellular and transmembrane domains of APP, thus its release implies the existence of

two distinct proteolytic events to generate its NH₂- and COOH-termini. At least two secretory mechanisms exist which release APP from the membrane and generate the soluble, COOH-truncated forms of APP (APP_s). Proteases which release APP and its fragments from the
5 membrane are termed "secretases". Most APP_s is released by a putative α-secretase which cleaves within the Aβ domain (between residues Lys¹⁶ and Leu¹⁷) to release α-APP_s and precludes the release of intact Aβ. A minor portion of APP_s is released by a β-secretase, which cleaves near the NH₂-terminus of Aβ and produces COOH-terminal fragments (CTFs) which
10 contain the whole Aβ domain. Finding these fragments in the extracellular compartment suggests that another proteolytic activity (γ-secretase) exists under normal conditions which can generate the COOH-terminus of Aβ.


It is believed that γ-secretase itself depends for its activity on the
15 presence of presenilin-1. In a manner that is not fully understood presenilin-1 appears to undergo autocleavage.

There are relatively few reports in the literature of compounds with inhibitory activity towards β- or γ-secretase, as measured in cell-based assays. These are reviewed in the articles referenced above. Many of the
20 relevant compounds are peptides or peptide derivatives.

WO 01/70677 discloses certain sulphonamido-substituted bridged bicycloalkyl derivatives which are useful in the treatment of Alzheimer's disease, but neither discloses nor suggests the compounds of the present invention.

25 The present invention provides a novel class of non-peptidic compounds which are useful in the treatment or prevention of AD by modulating the processing of APP by the putative γ-secretase, thus arresting the production of Aβ and preventing the formation of insoluble plaques.

30 According to the invention there is provided a compound of formula I:



I

wherein the moiety X-R is attached at one of the positions indicated by an asterisk;

X is a 5-(R-substituted)-1,2,4-oxadiazol-3-yl moiety; and

5 R is selected from:

(i) a non-aromatic hydrocarbon group of up to 10 carbon atoms, optionally substituted with halogen, CF₃, CHF₂, CN, OH, CO₂H, C₂₋₆acyl, C₁₋₄alkoxy or C₁₋₄alkoxycarbonyl;

10 (ii) a non-aromatic heterocyclic group comprising up to 7 ring atoms of which up to 3 are chosen from N, O and S and the remainder are carbon, bearing 0-3 substituents independently selected from oxo, halogen, CN, C₁₋₆alkyl, OH, CF₃, CHF₂, CH₂F, C₂₋₆acyl, CO₂H, C₁₋₄alkoxy and C₁₋₄alkoxycarbonyl; and

15 (iii) phenyl or 6-membered heteroaryl, either of which bears 0-3 substituents independently selected from halogen, CF₃, CHF₂, CH₂F, NO₂, CN, OCF₃, C₁₋₆alkyl and C₁₋₆alkoxy; or a pharmaceutically acceptable salt thereof.

In a preferred embodiment of the invention, R represents a group selected from those set forth under (iii) above.

20 It will be readily apparent to those skilled in the art that the compounds of formula I exist in two enantiomeric forms, depending on which of the ring positions indicated by an asterisk is bonded to the moiety -X-R. It is to be emphasised that the invention, for each identity of -X-R, encompasses both enantiomers, either as homochiral compounds or as mixtures of enantiomers in any proportion. Furthermore, structural formulae depicting attachment of -X-R or a synthetic precursor thereof at one of the said ring positions shall hereinafter be indicative of attachment

25

at either or both of said ring positions, unless expressly stated otherwise.

The homochiral compounds may be named as

[6S,9R,11R]2',3',4',5,5',6,7,8,9,10-decahydro-2-(5-(R-substituted)-1,2,4-oxadiazol-3-yl)-5'-(2,2,2-trifluoroethyl)spiro[6,9-methanobenzocyclooctene-11,3'-[1,2,5]thiadiazole] 1',1'-dioxides and [6R,9S,11S]2',3',4',5,5',6,7,8,9,10-decahydro-2-(5-(R-substituted)-1,2,4-oxadiazol-3-yl)-5'-(2,2,2-trifluoroethyl)spiro[6,9-methanobenzocyclooctene-11,3'-[1,2,5]thiadiazole] 1',1'-dioxides. The [6S,9R,11R] isomers are preferred.

Where a variable occurs more than once in formula I or in a substituent thereof, the individual occurrences of that variable are independent of each other, unless otherwise specified.

As used herein, the expression "C_{1-x}alkyl" where x is an integer greater than 1 refers to straight-chained and branched alkyl groups wherein the number of constituent carbon atoms is in the range 1 to x. Particular alkyl groups are methyl, ethyl, n-propyl, isopropyl and t-butyl. Derived expressions such as "C₂₋₆alkenyl", "hydroxyC₁₋₆alkyl", "heteroarylC₁₋₆alkyl", "C₂₋₆alkynyl" and "C₁₋₆alkoxy" are to be construed in an analogous manner. Most suitably, the number of carbon atoms in such groups is not more than 4.

The expression "C₃₋₆cycloalkyl" as used herein refers to nonaromatic monocyclic hydrocarbon ring systems comprising from 3 to 6 ring atoms. Examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cyclohexenyl.

The expression "C₃₋₆ cycloalkyl(C₁₋₆)alkyl" as used herein includes cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl.

The expression "C₂₋₆acyl" as used herein refers to (C₁₋₅alkyl)carbonyl groups, such as acetyl, propanoyl and butanoyl, including cycloalkyl derivatives such as cyclopentanecarbonyl and cyclobutanecarbonyl, and fluorinated derivatives such as trifluoroacetyl.

The term "halogen" as used herein includes fluorine, chlorine, bromine and iodine, of which fluorine and chlorine are preferred.

For use in medicine, the compounds of formula I may advantageously be in the form of pharmaceutically acceptable salts. Other salts may, however, be useful in the preparation of the compounds of formula I or of their pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds of this invention include acid addition salts which may, for example, be formed by mixing a solution of the compound according to the invention with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulphuric acid, methanesulphonic acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, oxalic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid. Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, e.g. sodium or potassium salts; alkaline earth metal salts, e.g. calcium or magnesium salts; and salts formed with suitable organic ligands, e.g. quaternary ammonium salts.

Where the compounds according to the invention have at least one asymmetric centre, they may accordingly exist as enantiomers. Where the compounds according to the invention possess two or more asymmetric centres, they may additionally exist as diastereoisomers. It is to be understood that all such isomers and mixtures thereof in any proportion are encompassed within the scope of the present invention.

In the compounds of formula I, X is a 5-(R-substituted)-1,2,4-oxadiazol-3-yl moiety.

In one embodiment, R is a non-aromatic hydrocarbon group of up to 10 carbon atoms, optionally substituted with halogen, CF₃, CHF₂, CN, OH, CO₂H, C₂₋₆acyl, C₁₋₄alkoxy or C₁₋₄alkoxycarbonyl. Within this definition, R may comprise cyclic or acyclic hydrocarbon residues or combinations thereof, saturated or unsaturated, up to a maximum of 10 carbon atoms in total, and may bear a substituent as detailed above. Within this


embodiment, R typically contains up to 6 carbon atoms. Suitable examples include alkyl groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, t-butyl and 2-methylpropyl, cycloalkyl groups such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl, cycloalkylalkyl groups such as cyclopropylmethyl and cyclopentylmethyl, alkenyl groups such as allyl, cyclopentenyl and cyclohexenyl, and alkynyl groups such as propargyl.

In a second embodiment, R represents a non-aromatic heterocyclic group comprising up to 7 ring atoms of which up to 3 are chosen from N, O and S and the remainder are carbon, bearing 0-3 substituents independently selected from oxo, halogen, CN, C₁₋₆alkyl, OH, CF₃, CHF₂, CH₂F, C₂₋₆acyl, CO₂H, C₁₋₄alkoxy and C₁₋₄alkoxycarbonyl. Suitable heterocyclic groups include pyrrolidine, piperidine, tetrahydropyridine, piperazine, morpholine, thiomorpholine, thiomorpholine-1,1-dioxide, pyran and thiopyran. Heterocyclic groups containing one or more nitrogen atoms may be bonded to X via carbon or via nitrogen. Within this embodiment, R very aptly represents 1-acetylpiperidin-4-yl.

In a third embodiment, R represents phenyl or 6-membered heteroaryl, either of which bears 0-3 substituents independently selected from halogen, CF₃, CHF₂, CH₂F, NO₂, CN, OCF₃, C₁₋₆alkyl and C₁₋₆alkoxy. Examples of suitable 6-membered heteroaryl groups represented by X include pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl and triazinyl. Preferably, the phenyl or heteroaryl ring bears 0 to 2 substituents. Phenyl groups represented by R preferably bear at least one substituent. Preferred substituents include halogen (especially chlorine and fluorine) and CF₃. Within this embodiment, examples of groups represented by R include monohalophenyl, dihalophenyl, pyridyl, monohalopyridyl and trifluoromethylpyridyl, wherein "halo" refers to fluoro or chloro. Suitable specific values for R include 2-fluorophenyl, 2-chlorophenyl, 4-fluorophenyl, 4-chlorophenyl, 2,4-difluorophenyl, 2,4-dichlorophenyl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, pyrazin-2-yl, 5-fluoropyridin-2-yl, 5-

chloropyridin-2-yl, 5-(trifluoromethyl)pyridin-2-yl and 6-(trifluoromethyl)pyridin-3-yl, and preferred examples include 4-fluorophenyl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, pyrazin-2-yl, 5-fluoropyridin-2-yl, 5-chloropyridin-2-yl, 5-(trifluoromethyl)pyridin-2-yl, and 6-(trifluoromethyl)pyridin-3-yl.

Representative compounds in accordance with the invention include those in which the moiety -X-R is selected from:



The compounds of the present invention have an activity as inhibitors of γ secretase.

The invention also provides pharmaceutical compositions comprising one or more compounds of this invention and a pharmaceutically acceptable carrier. Preferably these compositions are in unit dosage forms such as tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, transdermal patches, auto-injector devices or suppositories; for oral, parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums or surfactants such as sorbitan monooleate, polyethylene glycol, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is

meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules.

This solid preformulation composition is then subdivided into unit dosage
5 forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. Typical unit dosage forms contain from 1 to 100 mg, for example 1, 2, 5, 10, 25, 50 or 100 mg, of the active ingredient. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the
10 advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the
15 duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The present invention also provides a compound of the invention or
20 a pharmaceutically acceptable salt thereof for use in a method of treatment of the human body. Preferably the treatment is for a condition associated with the deposition of β -amyloid. Preferably the condition is a neurological disease having associated β -amyloid deposition such as Alzheimer's disease.

25 The present invention further provides the use of a compound of the present invention or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating or preventing Alzheimer's disease.

Also disclosed is a method of treatment of a subject suffering from or
30 prone to Alzheimer's disease which comprises administering to that


subject an effective amount of a compound according to the present invention or a pharmaceutically acceptable salt thereof.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection
5 include aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions include synthetic and natural gums such as
10 tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, poly(vinylpyrrolidone) or gelatin.

For treating or preventing Alzheimer's Disease, a suitable dosage level is about 0.01 to 250 mg/kg per day, preferably about 0.01 to 100 mg/kg per day, and especially about 0.01 to 50 mg/kg of body weight per
15 day. The compounds may be administered on a regimen of 1 to 4 times per day. In some cases, however, a dosage outside these limits may be used.


The compounds of the invention are particularly suitable for oral administration.

Suitable starting materials for the preparation of the compounds of
20 formula I are the ketones II:



II


wherein Y represents alkoxycarbonyl (especially CO₂Me or CO₂Et) or benzyloxy. Treatment of the ketones II with t-butylsulphonamide in the presence of TiCl₄ and triethylamine in refluxing dichloroethane provides
25 the sulphonylimines III(a):



III (a) $n = 2$
(b) $n = 1$

where Y has the same meaning as before. Alternatively, treatment of the ketones II with t-butylsulphinamide in the presence of $\text{Ti}(\text{OEt})_4$ in refluxing tetrahydrofuran provides the sulphonylimines III(b).


- 5 Both types of imine III react with trimethylsulphoxonium iodide to provide aziridines IV:



IV (a) $n = 2$
(b) $n = 1$

- where Y has the same meaning as before. The reaction takes place in the presence of sodium hydride at ambient temperature in a THF-DMSO mixture or at 0°C in DMSO.

Reaction of sulphonamides IV(a) with $\text{CF}_3\text{CH}_2\text{NH}_2$, followed by cleavage of the t-butylsulphonyl group, provides diamines V:




V

- where Y has the same meaning as before. Ring-opening of the aziridine is typically effected by heating with $\text{CF}_3\text{CH}_2\text{NH}_2$ at 100°C in DMF or DMSO in a sealed tube, while cleavage of the t-butylsulphonyl group may be effected by treatment with trifluoromethanesulphonic acid at 0°C .


- Alternatively, the diamines V may be formed directly by reaction of the sulphinamides IV(b) with $\text{CF}_3\text{CH}_2\text{NH}_2$ in the presence of zinc iodide in dichloroethane at 100°C .

Reaction of diamines V with sulphamide ($\text{H}_2\text{NSO}_2\text{NH}_2$) in refluxing anhydrous pyridine provides the cyclic sulphamides VI(a) and (b):




Compounds VI(a) and (b) are key intermediates for the preparation of compounds of formula I since the substituent Y may be transformed into the moiety -X-R by various standard synthetic techniques.

Hydrolysis of the esters VI(a) (for example, using sodium hydroxide in aqueous THF at 60°C) provides the acid VII which serves as the precursor for suitable heteroaryl structures.




The acid VII may be converted to the corresponding nitrile IX, which also serves as precursor for heteroaryl derivatives in accordance with formula I.



The nitrile is readily formed by first converting the acid to the corresponding carboxamide, then reacting it with triflic anhydride in the presence of triethylamine in dichloromethane at ambient temperature. The amide may be formed by treatment of the acid firstly with pentafluorophenol and dicyclohexylcarbodiimide, and then with ammonia.


The nitrile IX may be converted to the corresponding amidoxime by treatment with hydroxylamine hydrochloride in refluxing ethanolic sodium hydroxide solution. Subsequent coupling with a carboxylic acid RCO_2H and cyclisation provides a 1,2,4-oxadiazol-3-yl derivative X:



X

where R has the same meaning as before. The coupling may be carried out using carbonyldiimidazole in DMF at 70°C , while the final cyclisation may be effected by treatment with toluenesulphonic acid in a refluxing DMF-toluene mixture.


The benzyl ether VI(b) may be converted to the corresponding phenol by hydrogenation over Pd/C, and thence to the triflate XI by treatment with triflic anhydride in the presence of pyridine in dichloromethane at 0°C :



XI

where Tf represents triflyl (i.e. trifluoromethanesulphonyl). The triflate XI is another precursor of compounds in accordance with formula I.

Thus, XI may be converted to a boronic acid derivative XII:



XII

where R^2 represents H or alkyl, or the two OR^2 groups complete a cyclic boronate ester such as the pinacolate. The conversion may be achieved by conventional means using a suitable boron reagent, such as bis(pinacolato)diboron, in the presence of a Pd(II) catalyst such as
5 bis(diphenylphosphinoferrocene)dichloropalladium(II), typically in the presence of potassium acetate in DMF at 100°C.

Reaction of boronates XII with R-X-L, where L is a suitable leaving group (such as halogen, especially bromine or iodine) and R and X have the same meanings as before, provides compounds of formula I directly.
10 The reaction takes place in the presence of the same Pd catalyst as used in the preparation of XII.

Alternatively, the R-X moiety may be assembled in a three-stage process in which firstly an HX- group is introduced by reaction of boronate XII with HX-L, where L and X have the same meanings as before;
15 secondly, the resulting product is brominated to convert the HX- group to Br-X-; and thirdly, reaction with R-B(OR^2) provides a compound of formula I, where R and R^2 have the same meanings as before.

It will be apparent to those skilled in the art that, in many cases, the steps in the synthetic schemes described above may be carried out in a different order. Thus, if desired, the group Y in ketones II may be
20 converted to the moiety R-X- by one of the described routes prior to the construction of the spiro-linked sulphamide ring.

It will also be appreciated that where more than one isomer can be obtained from a reaction then the resulting mixture of isomers can be
25 separated by conventional means.

Where the above-described process for the preparation of the compounds according to the invention gives rise to mixtures of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The novel compounds may be
30 prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The novel compounds

may, for example, be resolved into their component enantiomers by standard techniques such as preparative HPLC, or the formation of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-p-toluoyl-d-tartaric acid and/or (+)-di-p-toluoyl-l-tartaric acid, followed by fractional crystallization and regeneration of the free base. The novel compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary.

In a preferred route to enantiomerically pure compounds of formula I, racemic intermediates VI(a) or VI(b) are subjected to preparative chiral HPLC to provide the corresponding homochiral intermediates, which are then converted to homochiral compounds of formula I by the routes indicated above.

Where they are not commercially available, the starting materials and reagents used in the above-described synthetic schemes may be prepared by conventional means. The ketones II may be prepared by the method described in *J.Org.Chem.*, **47**, 4329-34, 1987.

During any of the above synthetic sequences it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in *Protective Groups in Organic Chemistry*, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, *Protective Groups in Organic Synthesis*, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

A typical assay which can be used to determine the level of activity of compounds of the present invention is as follows:

(1) Mouse neuroblastoma neuro 2a cells expressing human app695 are cultured at 50-70% confluency in the presence of sterile 10mM sodium butyrate.

- (2) Cells are placed in 96-well plates at 30,000/well/100 μ L in minimal essential medium (MEM) (phenol red-free) + 10% foetal bovine serum (FBS), 50mM HEPES buffer (pH7.3), 1% glutamine, 0.2mg/ml G418 antibiotic, 10mM sodium butyrate.
- 5 (3) Make dilutions of the compound plate. Dilute stock solution to 5.5% DMSO/110 μ M compound. Mix compounds vigorously and store at 4°C until use.
- (4) Add 10 μ L compound/well. Mix plate briefly, and leave for 18h in 37°C incubator.
- 10 (5) Remove 90 μ L of culture supernatant and dilute 1:1 with ice-cold 25mM HEPES (pH.3), 0.1% BSA, 1.0mM EDTA (+ broad spectrum protease inhibitor cocktail; pre-aliquotted into a 96-well plate). Mix and keep on ice or freeze at -80°C.
- (6) Add back 100 μ L of warm MEM + 10% FBS, 50mM HEPES (pH7.3),
15 1% glutamine, 0.2mg/ml G418, 10mM sodium butyrate to each well, and return plate to 37°C incubator.
- (7) Prepare reagents necessary to determine amyloid peptide levels, for example by ELISA assay.
- (8) To determine if compounds are cytotoxic, cell viability following
20 compound administration is assessed by the use of redox dye reduction. A typical example is a combination of redox dye MTS (Promega) and the electron coupling reagent PES. This mixture is made up according to the manufacturer's instructions and left at room temperature.
- (9) Quantitate amyloid beta 40 and 42 peptides using an appropriate
25 volume of diluted culture medium by standard ELISA techniques.
- (10) Add 15 μ L/well MTS/PES solution to the cells; mix and leave at 37°C.
- (11) Read plate when the absorbance values are approximately 1.0 (mix briefly before reading to disperse the reduced formazan product).
- 30 Alternative assays are described in *Biochemistry*, 2000, 39(30), 8698-8704.


The compounds of the present invention show unexpectedly high affinities as measured by the above assays. Thus the following Examples all had an ED₅₀ of less than 50nM, typically less than 10nM, and frequently about 1nM in at least one of the above assays. In general, the compounds also exhibit good oral bioavailability and/or brain penetration, and are largely free from undesirable biological interactions with potential to cause toxic side effects.

The following examples illustrate the present invention.


EXAMPLES

Intermediate A

[6S/R,9R/S,11R/S] 2',3',4',5,5',6,7,8,9,10-Decahydro-2-carboxy-5'-(2,2,2-trifluoroethyl)-spiro[6,9-methanobenzocyclooctene-11,3'-[1,2,5]thiadiazole] 1',1'-dioxide




Step 1: Methyl 11-Oxo-5,6,7,8,9,10-hexahydro-6,9-methanobenzocyclooctene-2-carboxylate




Prepared using the procedure described for 11-oxo-5,6,7,8,9,10-hexahydro-6,9-methanobenzocyclooctene (Justus Liebigs Ann. Chem. **1961**, 650, 115) using methyl 3,4-bis(bromomethyl)benzoate in place of 1,2-bis(bromomethyl)benzene.

Step 2: Methyl 11-(2'-Methyl-propane-2'-sulfonylimino)-5,6,7,8,9,10-hexahydro-6,9-methanobenzocyclooctene-2-carboxylate




A solution of titanium (IV) chloride (25 mL, 1M in DCM) was added at r.t. to a stirred suspension of the ketone from Step 1 (10.0 g), *tert*-butyl sulfonamide (5.8 g) and triethylamine (11.6 mL) in 1,2-dichloroethane (100 mL). The yellow suspension was stirred under nitrogen for 30 minutes, then refluxed for 8 hours. The mixture was diluted with dichloromethane (400 mL) and saturated aqueous sodium hydrogencarbonate (400 mL) and filtered through a pad of Celite. The organic phase was dried (Na₂SO₄), filtered and concentrated to give the sulfonimine (15.5 g, quantitative) as a brown gum. δ (¹H, 360MHz, CDCl₃) 1.20-1.40 (2H, m), 1.50 (9H, s), 1.75-2.00 (2H, m), 2.90-2.95 (2H, m), 3.05-3.20 (3H, m), 3.92 (3H, s), 3.95-4.05 (1H, m), 7.24-7.28 (1H, m), 7.84-7.87 (2H, m).

Step 3: Methyl [6S/R,9R/S,11R/S]-11-[(2'-Methyl-propane-2'-sulfonyl)-1'-aza-spiro[2.4]]-5,6,7,8,9,10-hexahydro-6,9-methanobenzocyclooctene-2-carboxylate




Sodium hydride (55%, 1.75 g) was added portionwise at room temperature to a stirred suspension of trimethylsulfoxonium iodide (8.8 g) in DMSO (70 mL) under nitrogen. After stirring for 2 hours, the solution was cooled in an ice-bath and a solution of the sulfonimine from Step 2 (10.0 g) in DMSO (80 mL) was added. The mixture was stirred at room temperature for 2.5 hours then poured onto ice (500 mL) and diluted with water (200 mL). The white solids were collected, rinsed with diethyl ether and dried under vacuum at room temperature to give the aziridine (8.16 g, 79%). MS (ES+) 400 ([MNa]⁺).

Step 4: Methyl [6S/R,9R/S,11R/S]-11-(2'-Methyl-propane-2'-sulfonylamino)-11-(2,2,2-trifluoroethyl)aminomethyl-5,6,7,8,9,10-hexahydro-6,9-methanobenzocyclooctene-2-carboxylate




A solution of the aziridine from Step 3 (8.16 g) and 2,2,2-trifluoroethylamine (10.2 mL) in DMSO (45 mL) was divided into 3 equal portions and stirred at 100°C in sealed tubes for 6 days. The cooled solutions were combined and poured into water (250 mL). The white crystalline solid was collected, rinsed with diethyl ether and dried under vacuum at room temperature to give the amine (7.43 g, 72%). MS (ES+) 477 ([MH]⁺).

10 Step 5: Methyl [6S/R,9R/S,11R/S] 11-Amino-11-(2,2,2-trifluoroethyl)aminomethyl-5,6,7,8,9,10-hexahydro-6,9-methanobenzocyclooctene-2-carboxylate




Trifluoromethanesulfonic acid (11 mL) was added dropwise at 0°C to a stirred suspension of the sulfonamide from Step 4 (7.43 g) in dichloromethane (100 mL) under nitrogen. The mixture was stirred at 0°C for 20 minutes, then at room temperature for 4 hours. The mixture was poured into saturated aqueous sodium hydrogencarbonate (600 mL) and extracted with 10% methanol-dichloromethane (2x 300 mL). The extracts were washed with brine (100 mL), dried over Na₂SO₄, filtered and concentrated to give the diamine (6.5 g, quantitative) as a yellow oil. MS (ES+) 357 ([MH]⁺).

25 Step 6: [6S/R,9R/S,11R/S] 2',3',4',5,5',6,7,8,9,10-Decahydro-2-carbomethoxy-5'-(2,2,2-trifluoroethyl)-spiro[6,9-methanobenzocyclooctene-11,3'-[1,2,5]thiadiazole] 1',1'-dioxide



A solution of the diamine from Step 5 (6.5 g) and sulfamide (4.5 g) was relaxed in pyridine (120 mL) for 6 hours. Solvent was removed by evaporation. The residue was partitioned between 1M HCl (200 mL) and
 5 10% methanol-dichloromethane (500 mL). The organic extract was washed with brine (100 mL), dried over Na₂SO₄, filtered and concentrated. The solid was rinsed with diethyl ether to give the cyclic sulfamide (5.26 g, 80%) as an off-white powder. δ (¹H, 360MHz, CDCl₃) 1.26-1.30 (2H, m), 1.70-1.75 (2H, m), 2.47-2.50 (2H, m), 2.74-2.82 (2H, m), 3.21-3.29 (2H, m),
 10 3.41-3.47 (2H, m), 3.68 (2H, q, *J*=9), 3.90 (3H, s), 4.78 (1H, s), 7.17 (1H, d, *J*=8), 7.78-7.80 (2H, m)


Step 7: [6S/R,9R/S,11R/S] 2',3',4',5,5',6,7,8,9,10-Decahydro-2-carboxy-5'-(2,2,2-trifluoroethyl)-spiro[6,9-methanobenzocyclooctene-11,3'-[1,2,5]thiadiazole] 1',1'-dioxide



A mixture of 4M sodium hydroxide (3 mL) and the ester from Step 6 (2.33 g) in tetrahydrofuran (30 mL) was stirred at 60°C for 3.5 hours. The mixture was diluted with 1M citric acid (10 mL) and 1M hydrochloric acid
 20 (20 mL) and extracted with dichloromethane (2 x 100 mL). The extracts were washed with water (100 mL), brine (100 mL), dried over Na₂SO₄, filtered and concentrated. Trituration with diethyl ether – isohexane gave the acid (1.96 g, 88%) as an off-white solid. δ (¹H, 360MHz, d₆-DMSO) 0.95-1.05 (2H, m), 1.65-1.75 (2H, m), 2.35-2.42 (2H, m), 2.62-2.72 (2H, m), 3.18-3.23 (2H, m), 3.47 (2H, s), 4.02 (2H, q, *J*=10), 7.23 (1H, d, *J*=8), 7.62-7.70
 25 (2H, m), 8.03 (1H, s), 12.60 (1H, s).


Example 1

[6S/R,9R/S,11R/S]-2',3',4',5,5',6,7,8,9,10-Decahydro-2-(5-(4-fluorophenyl)-1,2,4-oxadiazol-3-yl)-5'-(2,2,2-trifluoroethyl)spiro[6,9-methanobenzocyclooctene-11,3'-[1,2,5]thiadiazole] 1',1'-dioxide



5

Step 1 : [6S/R,9R/S,11R/S]-2',3',4',5,5',6,7,8,9,10-Decahydro-2-amidoxime-5'-(2,2,2-trifluoroethyl)spiro[6,9-methanobenzocyclooctene-11,3'-[1,2,5]thiadiazole] 1',1'-dioxide



- 10 Pentafluorophenol (0.341 g, 1.85 mmol) and *N,N'*-dicyclohexylcarbodiimide (0.382 g, 1.85 mmol) were added to a suspension of Intermediate A (0.5 g, 1.24 mmol) in ethyl acetate (30 mL) and the reaction was stirred at room temperature for 1 hour after which time it was homogeneous. The solvent was removed *in vacuo*. Ethyl acetate was added to the residue and the
- 15 resulting white solid removed by filtration. The filtrate was evaporated *in vacuo*. The residue was dissolved in dichloromethane (50 ml) and ammonia gas was bubbled through until the mixture was cloudy. The reaction was stirred for 1 hour at room temperature. The solvent was removed *in vacuo* and the residue purified by flash column
- 20 chromatography on silica, eluting with 20% ethyl acetate – isohexane followed by 5% methanol – dichloromethane, to give the carboxamide (0.385 g, 77%).


A solution of this (0.2 g, 0.5 mmol) and (methoxycarbonylsulfamoyl)triethyl-ammonium hydroxide, inner salt,

25 (0.177 g, 1.74 mmol) in tetrahydrofuran (10 mL) was heated under reflux

for 4 hours. The solvent was removed *in vacuo* and the residue purified by flash column chromatography on silica, eluting with 25% ethyl acetate – isohexane, to give the nitrile (0.115 g, 60%).

A solution of this (0.1 g, 0.26 mmol), hydroxylamine hydrochloride (0.027 g, 0.39 mmol) and triethylamine (65 μ L, 0.47 mmol) in ethanol (5 mL) was heated under reflux for 18 hours. The cooled reaction mixture was poured into sodium hydroxide solution (1M aq; 25 mL) and extracted with ethyl acetate (25 mL). The organic layer was washed with water (25 mL) and brine (25 mL), dried over Na₂SO₄ and evaporated to give the amidoxime (0.038 g, 35%). MS (ES+) 419, 420, 421.

Step 2 : [6S/R,9R/S,11R/S]-2',3',4',5,5',6,7,8,9,10-Decahydro-2-(5-(4-fluorophenyl)-1,2,4-oxadiazol-3-yl)-5'-(2,2,2-trifluoroethyl)spiro[6,9-methanobenzocyclooctene-11,3'-[1,2,5]thiadiazole] 1',1'-dioxide




purified by flash column chromatography on silica, eluting with 25% ethyl acetate – isohexane to give the title compound (0.055 g, 48%).

5 δ (^1H , 400MHz, d_4 -MeOH) 1.22-1.30 (2H, m), 1.78-1.81 (2H, m), 2.48-2.53 (2H, m), 2.73-2.82 (2H, m), 3.39 (2H, dd, $J=5, 16$), 3.52 (2H, s), 3.88 (2H, q, $J=9$), 7.29 (1H, d, $J=7.8$), 7.34-7.40 (2H, m), 7.86-7.89 (2H, m), 8.26-8.30 (2H, m); MS (ES+) 523, 524, 526 ($[\text{MH}]^+$).

Example 2

10 **[6S/R,9R/S,11R/S] 2',3',4',5,5',6,7,8,9,10-Decahydro-2-(5-(2-pyridyl)-1,2,4-oxadiazol-3-yl)-5'-(2,2,2-trifluoroethyl)spiro[6,9-methanobenzocyclooctene-11,3'-[1,2,5]thiadiazole] 1',1'-dioxide**




Prepared in the same way as Example 1 using 0.090 g of amidoxime from
15 Step 1 and picolinic acid (0.024 g, 0.19 mmol) to give the title compound
(0.046 g, 41%).

MS (ES+) 506, 507, 508.

Example 3

20 **[6S,9R,11R] 2',3',4',5,5',6,7,8,9,10-Decahydro-2-(5-(pyridin-2-yl)-1,2,4-oxadiazol-3-yl)-5'-(2,2,2-trifluoroethyl)-spiro[6,9-methanobenzocyclooctene-11,3'-[1,2,5]thiadiazole] 1',1'-dioxide**




25 The racemic oxadiazole Example 2 was resolved by supercritical fluid chromatographic separation on a Chiralcel OD column using 30%

methanol - sc-CO₂ as eluant (50 mL/min) at 35°C, 100 bar and UV detection at 250 nm, collecting the second compound eluted under these conditions.

5 Example 4

[6S/R,9R/S,11R/S]-2',3',4',5,5',6,7,8,9,10-Decahydro-2-(5-(3-pyridyl)-1,2,4-oxadiazol-3-yl)-5'-(2,2,2-trifluoroethyl)spiro[6,9-methanobenzocyclooctene-11,3'-[1,2,5]thiadiazole] 1',1'-dioxide




Prepared in the same way as Example 1 using 0.14 g of amidoxime from step 1 and nicotinic acid (0.037 g, 0.3 mmol) to give the title compound (0.027 g, 16%).

MS (ES+) 506, 507, 508, 547, 548

15

CLAIMS

1. A compound of formula I:



I

wherein the moiety X-R is attached at one of the positions indicated by an asterisk;

X is a 5-(R-substituted)-1,2,4-oxadiazol-3-yl moiety; and

R is selected from:

- (i) a non-aromatic hydrocarbon group of up to 10 carbon atoms, optionally substituted with halogen, CF₃, CHF₂, CN, OH, CO₂H, C₂₋₆acyl, C₁₋₄alkoxy or C₁₋₄alkoxycarbonyl;
 - (ii) a non-aromatic heterocyclic group comprising up to 7 ring atoms of which up to 3 are chosen from N, O and S and the remainder are carbon, bearing 0-3 substituents independently selected from oxo, halogen, CN, C₁₋₆alkyl, OH, CF₃, CHF₂, CH₂F, C₂₋₆acyl, CO₂H, C₁₋₄alkoxy and C₁₋₄alkoxycarbonyl; and
 - (iii) phenyl or 6-membered heteroaryl, either of which bears 0-3 substituents independently selected from halogen, CF₃, CHF₂, CH₂F, NO₂, CN, OCF₃, C₁₋₆alkyl and C₁₋₆alkoxy;
- or a pharmaceutically acceptable salt thereof.


2. A compound according to claim 1 which is homochiral and is [6S,9R,11R]2',3',4',5,5',6,7,8,9,10-decahydro-2-(5-(R-substituted)-1,2,4-oxadiazol-3-yl)-5'-(2,2,2-trifluoroethyl)spiro[6,9-methanobenzocyclooctene-11,3'-[1,2,5]thiadiazole] 1',1'-dioxide, or a pharmaceutically acceptable salt thereof.

3. A compound according to claim 1 or claim 2 wherein R represents phenyl or a 6-membered heteroaryl, either of which bears 0-3 substituents independently selected from halogen, CF₃, CHF₂, CH₂F, NO₂, CN, OCF₃,
5 C₁₋₆alkyl and C₁₋₆alkoxy.

4. A compound according to claim 3 wherein R is selected from monohalophenyl, dihalophenyl, pyridyl, monohalopyridyl and trifluoromethylpyridyl, wherein "halo" refers to fluoro or chloro.

10

5. A compound according to claim 1 or claim 2 wherein the moiety -X-R is selected from:



15

6. A pharmaceutical composition comprising a compound according to any previous claim and a pharmaceutically acceptable carrier.

7. A compound according to any of claims 1-5 for use in a method of
20 treatment of the human body.

8. The use of a compound according to any of claims 1-5 in the manufacture of a medicament for treating or preventing Alzheimer's disease.

25

9. A method of treatment of a subject suffering from or prone to Alzheimer's disease which comprises administering to that subject an effective amount of a compound according to any of claims 1-5.

INTERNATIONAL SEARCH REPORT

PCT/GB 03/01777

A. CLASSIFICATION OF SUBJECT MATTER		
IPC 7 C07D417/04 C07D417/14 A61K31/433 A61K31/4439 A61P25/28		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07D A61K A61P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 01 70677 A (MERCK SHARP & DOHME LIMITED ET AL) 27 September 2001 (2001-09-27) cited in the application the whole document, particularly examples 177, 178, 200 and 201 ---	1,5-9
P,X	WO 02 36555 A (MERCK SHARP & DOHME LIMITED) 10 May 2002 (2002-05-10) the whole document, particularly examples 140-142 -----	1-9
<input type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *&* document member of the same patent family		
Date of the actual completion of the international search 11 July 2003		Date of mailing of the international search report 23/07/2003
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Allard, M

INTERNATIONAL SEARCH REPORT

PCT/GB 03/01777

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claim 9 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

PCT/GB 03/01777

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0170677	A	27-09-2001	AU 4086101 A	03-10-2001
			CA 2404125 A1	27-09-2001
			EP 1268412 A1	02-01-2003
			WO 0170677 A1	27-09-2001
			AU 1074702 A	15-05-2002
			WO 0236555 A1	10-05-2002
WO 0236555	A	10-05-2002	AU 1074702 A	15-05-2002
			AU 4086101 A	03-10-2001
			CA 2404125 A1	27-09-2001
			EP 1268412 A1	02-01-2003
			WO 0170677 A1	27-09-2001
			WO 0236555 A1	10-05-2002